Effective Date: 08/01/2021



NHS MEDICAL POLICY

Casirivimab and Imdevimab for COVID -19 + Patients Medicine 2021-001

A. May be indicated when ALL the following are present:

- 1 Administered together for the treatment of mild to moderate COVID-19 patients
- Given to Adults and pediatric patients (12 years of age or older weighing at least 40 kilograms [about 88 pounds]
- Positive results of direct SARS-Co-V-2 viral testing and who are at high risk for progressing to severe COVID-19.
- 4 Must have at least one of the following co-morbidities:
 - Cancer
 - Chronic kidney disease
 - Chronic lung disease (COPD, asthma [moderate to severe]. Interstitial lung disease, cystic fibrosis, and pulmonary hypertension)
 - Dementia or other neurological condition
 - Diabetes (type 1 or type 2)
 - Down syndrome
 - Heart conditions (such as heart failure, coronary artery disease, cardiomyopathies, or hypertension
 - HIV infection
 - Immunocompromised state
 - Liver disease such as alcohol-related liver disease, nonalcoholic fatty liver disease and cirrhosis
 - Overweight and obesity (BMI >25)
 - Pregnancy and recently pregnant people (for at least 42 days following end of pregnancy)
 - Sickle cell disease or thalassemia
 - Smoking, current or former
 - Solid organ or blood stem cell transplant
 - Stroke or cerebrovascular disease
 - Substance abuse disorders
- 5 Not hospitalized due to COVID-19

Does not require oxygen

SOURCES

U. S. Food and Drug Administration Emergency Use Authorization

CODE REFERENCE: (This may not be a comprehensive list of codes to apply to this policy.)

Q0243, Q0244, M0243, M0244

POLICY HISTORY/REVISION INFORMATION

Date	Action/Description
09/19/2022	Annual review and approval by UM Committee
08/23/2023	Annual review and approval by UM/QM Committee